

Uniwersytet Wrocławski
Wydział Matematyki i Informatyki
Instytut Matematyczny
specjalność: ISIM

Franciszek Malinka

Generic automorphisms as Fraïssé limits

Praca licencjacka
napisana pod kierunkiem
dra Tomasza Rzepeckiego

Wrocław 2022

This page is intentionally left blank

1. INTRODUCTION

Model theory is a field of mathematics that classifies and constructs structures with particular properties (particularly those expressible in first order logic). It describes classical mathematical objects in a broader context, abstracts their properties and studies connections between seemingly unrelated structures. This work studies limits of Fraïssé classes with additional combinatorial and categorical properties. Fraïssé classes are frequently used in model theory, both as a source of examples and to analyse particular “generic” structures.

The notion of Fraïssé class and its limit is due to the French logician Roland Fraïssé. He also introduced the back-and-forth argument, a fundamental model theoretical method in construction of elementarily equivalent structures, upon which Ehrenfeucht-Fraïssé games are based.

The prototypical example for this paper is the random graph 3.13 (also known as the Rado graph), the Fraïssé limit of the class of finite undirected graphs. It serves as a useful example, gives an intuition of the Fraïssé limits, weak Hrushovski property and free amalgamation. Perhaps most importantly, the random graph has a so-called generic automorphism 2.6, which was first proved by Truss in [10], where he also introduced the term.

The key Theorem 4.5 says that a Fraïssé class with canonical amalgamation and weak Hrushovski property has a generic automorphism. The fact that such an automorphism exists in this case follows from the classical results of Ivanov [3] and Kechris-Rosendal [5]. In this work we show a new way to construct a generic automorphism by expanding the structures of the class by a (total) automorphism and considering limit of such extended Fraïssé class. We achieve this by using the Banach-Mazur games, a well known method in the descriptive set theory, which proves useful in the study of comeagre sets.

WSTĘP

Teoria modeli jest działem matematyki zajmującym się klasyfikacją i konstrukcją struktur z określonymi cechami (szczególnie takimi, które da się wyrazić logiką pierwszego rzędu). Opisuje klasyczne matematyczne obiekty w szerszym kontekście, abstrahuje ich własności i opisuje połączenia między pozornie niepowiązanymi strukturami. Niniejsza praca bada granicę klas Fraïsségo z dodatkowymi kombinatorycznymi i kategoryjnymi własnościami. Klasy Fraïsségo są powszechnie znanym i używanym konceptem w teorii modeli, zarówno jako narzędzie opisujące “generyczne” struktury, jak i źródło przykładów.

Klasy Fraïsségo i ich granice zostały opisane po raz pierwszy przez francuskiego logika Rolanda Fraïsségo. Zawdzięczamy mu również argument “back-and-forth”, fundamentalną teoriomodelową metodę konstrukcji elementarnie równoważnych struktur, na podstawie której bazują gry Ehrenfeuchta-Fraïsségo.

Graf losowy 3.13, zwany również grafem Rado, jest prototypową strukturą tej pracy. Graf losowy można skonstruować jako granicę Fraïsségo klasy skończonych grafów nieskierowanych. Służy on jako użyteczny przykład, daje intuicję stojącą za konstrukcją granicy Fraïsségo, słabej własności Hrushovskiego oraz wolnej amalgamacji. Ponadto, co najważniejsze dla niniejszej pracy, graf losowy ma tak zwany *generyczny automorfizm* 2.6, co zostało po raz pierwsze zdefiniowane i udowodnione przez Trussa w [10].

Kluczowe twierdzenie 4.5 mówi, że klasa Fraïsségo z kanoniczną amalgamacją i słabą własnością Hrushovskiego ma generyczny automorfizm. Istnienie takiego automorfizmu w tym przypadku wynika z wcześniejszych klasycznych wyników Ivanova [3] oraz Kechrisa-Rosendala [5]. W tej pracy pokazujemy nowy sposób konstrukcji generycznego automorfizmu poprzez rozszerzenie struktur klasy o (totalny) automorfizm oraz analizę granicy Fraïsségo nowo powstałej klasy. Posługujemy się przy tym gramami Banacha-Mazura, które są dobrze znanym narzędziem w deskryptywnej teorii mnogości.

2. PRELIMINARIES

Before we get to the main work of the paper, we need to establish basic notions, known facts and theorems. This section provides a brief introduction to the theory of Baire spaces and category theory. Most of the notions are well known, interested reader may look at [4], [6]

2.1. Descriptive set theory. In this section we provide an important definition of a *comeagre* set. It is purely topological notion, the intuition may come from the measure theory though. For example, in a standard Lebesgue measure on the real interval $[0, 1]$, the set of rationals is of measure 0, although being a dense subset of the $[0, 1]$. So, in a sense, the set of rationals is *meagre* in the interval $[0, 1]$. On the other hand, the set of irrational numbers is also dense, but have measure 1, so it is *comeagre*.

This is only a rough approximation of the topological definition. The definitions are based on the Kechris' book *Classical Descriptive Set Theory* [4]. One should look into it for more details and examples.

Definition 2.1. Suppose X is a topological space and $A \subseteq X$. We say that A is *meagre* in X if $A = \bigcup_{n \in \mathbb{N}} A_n$, where A_n are nowhere dense subsets of X (i.e. $\text{Int}(\bar{A}_n) = \emptyset$).

Definition 2.2. We say that A is *comeagre* in X if it is a complement of a meagre set. Equivalently, a set is comeagre if and only if it contains a countable intersection of open dense sets.

Every countable set is meagre in any T_1 space. So, \mathbb{Q} is meagre in \mathbb{R} (although it is dense), which means that the set of irrationals is comeagre. The Cantor set is nowhere dense, hence meagre in the $[0, 1]$ interval.

Definition 2.3. We say that a topological space X is a *Baire space* if every comeagre subset of X is dense in X (equivalently, every meagre set has empty interior).

Definition 2.4. Suppose X is a Baire space. We say that a property P holds *generically* for a point $x \in X$ if $\{x \in X \mid P \text{ holds for } x\}$ is comeagre in X .

Let M be a structure. We define a topology on the automorphism group $\text{Aut}(M)$ by the basis of open sets: for a finite function $f : M \rightarrow M$ we have a basic open set $[f]_{\text{Aut}(M)} = \{g \in \text{Aut}(M) \mid f \subseteq g\}$. This is a standard definition.

Fact 2.5. For a countable structure M , the topological space $\text{Aut}(M)$ is a Baire space.

This is in fact a very weak statement, it is also true that $\text{Aut}(M)$ is a Polish space (i.e. separable completely metrizable), and every Polish space is Baire. However, those additional properties are not important in this study.

Definition 2.6. Let $G = \text{Aut}(M)$ be the automorphism group of structure M . We say that $f \in G$ is a *generic automorphism*, if the conjugacy class of f is comeagre in G .

Definition 2.7. Let X be a nonempty topological space and let $A \subseteq X$. The *Banach-Mazur game* of A , denoted as $G^{**}(A)$ is defined as follows: Players I and II take turns in playing nonempty open sets $U_0, V_0, U_1, V_1, \dots$ such that $U_0 \supseteq V_0 \supseteq U_1 \supseteq V_1 \supseteq \dots$. We say that player II wins the game if $\bigcap_n V_n \subseteq A$.

There is an important Theorem 2.13 on the Banach-Mazur game: A is comeagre if and only if II can always choose sets V_0, V_1, \dots such that it wins. Before we prove it we need to define notions necessary to formalise and prove the theorem.

Definition 2.8. T is the tree of all legal positions in the Banach-Mazur game $G^{**}(A)$ when T consists of all finite sequences (W_0, W_1, \dots, W_n) , where W_i are nonempty open sets such that $W_0 \supseteq W_1 \supseteq \dots \supseteq W_n$.

Definition 2.9. We say that σ is a pruned subtree of the tree of all legal positions T if $\sigma \subseteq T$, for any $(W_0, W_1, \dots, W_n) \in \sigma, n \geq 0$ there is a W such that $(W_0, W_1, \dots, W_n, W) \in \sigma$ (it simply means that there's no finite branch in σ) and $(W_0, W_1, \dots, W_{n-1}) \in \sigma$ (every node on a branch is in σ).

Definition 2.10. Let σ be a pruned subtree of the tree of all legal positions T . By $[\sigma]$ we denote the set of all infinite branches of σ , i.e. infinite sequences (W_0, W_1, \dots) such that $(W_0, W_1, \dots, W_n) \in \sigma$ for any $n \in \mathbb{N}$.

Definition 2.11. A strategy for II in $G^{**}(A)$ is a pruned subtree $\sigma \subseteq T$ such that

- (i) σ is nonempty,
- (ii) if $(U_0, V_0, \dots, U_n, V_n) \in \sigma$, then for all open nonempty $U_{n+1} \subseteq V_n$, $(U_0, V_0, \dots, U_n, V_n, U_{n+1}) \in \sigma$,
- (iii) if $(U_0, V_0, \dots, U_n) \in \sigma$, then for a unique V_n , $(U_0, V_0, \dots, U_n, V_n) \in \sigma$.

Intuitively, a strategy σ works as follows: I starts playing U_0 as any open subset of X , then II plays unique (by (iii)) V_0 such that $(U_0, V_0) \in \sigma$. Then I responds by playing any $U_1 \subseteq V_0$ and II plays unique V_1 such that $(U_0, V_0, U_1, V_1) \in \sigma$, etc.

We will often denote a sequence $U_0 \supseteq V_0 \supseteq U_1 \supseteq V_1 \supseteq \dots$ of open sets as an instance of a Banach-Mazur game, or just simply by a game.

Definition 2.12. A strategy σ is a winning strategy for II if for any instance $(U_0, V_0, \dots) \in [\sigma]$ of the Banach-Mazur game player II wins, i.e. $\bigcap_n V_n \subseteq A$.

Theorem 2.13 (Banach-Mazur, Oxtoby). *Let X be a nonempty topological space and let $A \subseteq X$. Then A is comeagre $\Leftrightarrow II$ has a winning strategy in $G^{**}(A)$.*

The statement of the theorem is once again taken from Kechris [4] 8.33. However, the proof given in the book is brief, thus we present a detailed version. In order to prove the theorem we add an auxiliary definition and lemma.

Definition 2.14. Let $S \subseteq \sigma$ be a pruned subtree of tree of all legal positions T and let $p = (U_0, V_0, \dots, V_n) \in S$. We say that S is comprehensive for p if the family $\mathcal{V}_p = \{V_{n+1} \mid (U_0, V_0, \dots, V_n, U_{n+1}, V_{n+1}) \in S\}$ (it may be that $n = -1$, which means $p = \emptyset$) is pairwise disjoint and $\bigcup \mathcal{V}_p$ is dense in V_n (where we put $V_{-1} = X$). We say that S is comprehensive if it is comprehensive for each $p = (U_0, V_0, \dots, V_n) \in S$.

Fact 2.15. *If σ is a winning strategy for II then there exists a nonempty comprehensive $S \subseteq \sigma$.*

Proof. We construct S recursively as follows:

- (1) $\emptyset \in S$,
- (2) if $(U_0, V_0, \dots, U_n) \in S$, then $(U_0, V_0, \dots, U_n, V_n) \in S$ for the unique V_n given by the strategy σ ,
- (3) let $p = (U_0, V_0, \dots, V_n) \in S$. For a possible player move of player I $U_{n+1} \subseteq V_n$ let U_{n+1}^* be the unique set player II would respond with by σ . Now, by Zorn's Lemma, let \mathcal{U}_p be a maximal collection of nonempty open subsets $U_{n+1} \subseteq V_n$ such that the set $\{U_{n+1}^* \mid U_{n+1} \in \mathcal{U}_p\}$ is pairwise disjoint. Then put in S all $(U_0, V_0, \dots, V_n, U_{n+1})$ such that $U_{n+1} \in \mathcal{U}_p$. This way S is comprehensive for p : the family $\mathcal{V}_p = \{V_{n+1} \mid (U_0, V_0, \dots, V_n, U_{n+1}, V_{n+1}) \in S\}$ is exactly $\{U_{n+1}^* \mid U_{n+1} \in \mathcal{U}_p\}$, which is pairwise disjoint and $\bigcup \mathcal{V}_p$ is obviously dense in V_n by the maximality of \mathcal{U}_p – if there was any open set $\tilde{U}_{n+1} \subseteq V_n$ disjoint from $\bigcup \mathcal{V}_p$, then $\tilde{U}_{n+1}^* \subseteq \tilde{U}_{n+1}$ would be also disjoint from $\bigcup \mathcal{V}_p$, so the family $\mathcal{U}_p \cup \{\tilde{U}_{n+1}\}$ would violate the maximality of \mathcal{U}_p . \square

Lemma 2.16. *Let S be a nonempty comprehensive pruned subtree of a strategy σ . Then:*

- (i) *For any open $V_n \subseteq X$ there is at most one $p = (U_0, V_0, \dots, U_n, V_n) \in S$.*

Let $S_n = \{V_n \mid (U_0, V_0, \dots, V_n) \in S\}$ for $n \in \mathbb{N}$ (i.e. S_n is a family of all possible choices player II can make in its n -th move according to S).

- (ii) $\bigcup S_n$ *is open and dense in X .*
- (iii) S_n *is a family of pairwise disjoint sets.*

Proof. (i): Suppose that there are some $p = (U_0, V_0, \dots, U_n, V_n)$, $p' = (U'_0, V'_0, \dots, U'_n, V'_n)$ such that $V_n = V'_n$ and $p \neq p'$. Let k be the smallest index such that those sequences differ. We have two possibilities:

- $U_k = U'_k$ and $V_k \neq V'_k$ – this cannot be true simply by the fact that S is a subset of a strategy (so V_k is unique for U_k).
- $U_k \neq U'_k$: by the comprehensiveness of S we know that for $q = (U_0, V_0, \dots, U_{k-1}, V_{k-1})$ the set \mathcal{V}_q is pairwise disjoint. Thus $V_k \cap V'_k = \emptyset$, because $V_k, V'_k \in \mathcal{V}_q$. But this leads to a contradiction – V_n cannot be a nonempty subset of both V_k, V'_k .

(ii): The lemma is proved by induction on n . For $n = 0$ it follows trivially from the definition of comprehensiveness. Now suppose the lemma is true for n . Then the set $\bigcup_{V_n \in S_n} \bigcup \mathcal{V}_{p_{V_n}}$ (where p_{V_n} is given uniquely from (i)) is dense and open in X by the induction hypothesis. But $\bigcup S_{n+1}$ is exactly this set, thus it is dense and open in X .

(iii): We will prove it by induction on n . Once again, the case $n = 0$ follows from the comprehensiveness of S . Now suppose that the sets in S_n are pairwise disjoint. Take some $x \in V_{n+1} \in S_{n+1}$. Of course $\bigcup S_n \supseteq \bigcup S_{n+1}$, thus by the inductive hypothesis $x \in V_n$ for the unique $V_n \in S_n$. It must be that $V_{n+1} \in \mathcal{V}_{p_{V_n}}$, because V_n is the only superset of V_{n+1} in S_n . But $\mathcal{V}_{p_{V_n}}$ is disjoint, so there is no other $V'_{n+1} \in \mathcal{V}_{p_{V_n}}$ such that $x \in V'_{n+1}$. Moreover, there is no such set in $S_{n+1} \setminus \mathcal{V}_{p_{V_n}}$, because those sets are disjoint from V_n . Hence there is no $V'_{n+1} \in S_{n+1}$ other than V_n such that $x \in V'_{n+1}$. We have chosen x and V_{n+1} arbitrarily, so S_{n+1} is pairwise disjoint. \square

Now we can move to the proof of the Banach-Mazur theorem.

Proof of Theorem 2.13. \Rightarrow : Let (A_n) be a sequence of dense open sets with $\bigcap_n A_n \subseteq A$. Then II simply plays $V_n = U_n \cap A_n$, which is nonempty by the denseness of A_n .

\Leftarrow : Suppose II has a winning strategy σ . We will show that A is comeagre. Take a comprehensive $S \subseteq \sigma$. We claim that $\mathcal{S} = \bigcap_n \bigcup S_n \subseteq A$. By the lemma 2.16, (ii) sets $\bigcup S_n$ are open and dense, thus A must be comeagre. Now we prove the claim towards contradiction.

Suppose there is $x \in \mathcal{S} \setminus A$. By the lemma 2.16, (iii) for any n there is unique $x \in V_n \in S_n$. It follows that $p_{V_0} \subset p_{V_1} \subset \dots$. Now the game $(U_0, V_0, U_1, V_1, \dots) = \bigcup_n p_{V_n} \in [S] \subseteq [\sigma]$ is not winning for player II , which contradicts the assumption that σ is a winning strategy. \square

Corollary 2.17. *If we add a constraint to the Banach-Mazur game such that players can only choose basic open sets, then the Theorem 2.13 still suffices.*

Proof. If one adds the word *basic* before each occurrence of word *open* in previous proofs and theorems then they still will be valid (except for \Rightarrow , but its an easy fix – take for V_n a basic open subset of $U_n \cap A_n$). \square

This corollary will be important in using the theorem in practice – it’s much easier to work with basic open sets rather than arbitrary open sets.

2.2. Category theory. In this section we will give a short introduction to the notions of category theory that will be necessary to generalize the key result of the paper.

We will use a standard notation. If the reader is interested in a more detailed introduction to the category theory, then it’s recommended to take a look at [6]. Here we will shortly describe the standard notation.

A *category* \mathcal{C} consists of a collection of objects (denoted as $\text{Obj}(\mathcal{C})$, but most often simply as \mathcal{C}) and a collection of *morphisms* $\text{Mor}(A, B)$ between each pair of objects $A, B \in \mathcal{C}$. We require that for each pair of morphisms $f : B \rightarrow C$, $g : A \rightarrow B$ there was a morphism $f \circ g : A \rightarrow C$. If $f : A \rightarrow B$ then we say that A is the domain of f ($\text{Dom } f$) and that B is the range of f ($\text{Rng } f$).

For every $A \in \mathcal{C}$ there is an *identity morphism* $\text{id}_A : A \rightarrow A$ such that for any morphism $f \in \text{Mor}(A, B)$ we have that $f \circ \text{id}_A = \text{id}_B \circ f$.

We say that $f : A \rightarrow B$ is an *isomorphism* if there is (necessarily unique) morphism $g : B \rightarrow A$ such that $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$. Automorphism is an isomorphism where $A = B$.

A *functor* is a “(homo)morphism“ of categories. We say that $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor from category \mathcal{C} to category \mathcal{D} if it associates each object $A \in \mathcal{C}$ with an object $F(A) \in \mathcal{D}$, associates each morphism $f : A \rightarrow B$ in \mathcal{C} with a morphism $F(f) : F(A) \rightarrow F(B)$. We also require that $F(\text{id}_A) = \text{id}_{F(A)}$ and that for any (compatible) morphisms f, g in \mathcal{C} , $F(f \circ g) = F(f) \circ F(g)$ should hold.

In category theory we distinguish *covariant* and *contravariant* functors. Here, we only consider covariant functors, so we will simply say *functor*.

Fact 2.18. *Functor $F : \mathcal{C} \rightarrow \mathcal{D}$ maps isomorphism $f : A \rightarrow B$ in \mathcal{C} to the isomorphism $F(f) : F(A) \rightarrow F(B)$ in \mathcal{D} .*

A notion that will be very important for us is a “morphism of functors“ which is called *natural transformation*.

Definition 2.19. Let F, G be functors between the categories \mathcal{C}, \mathcal{D} . A *natural transformation* η is function that assigns to each object A of \mathcal{C} a morphism η_A in $\text{Mor}(F(A), G(A))$ such that for every morphism $f : A \rightarrow B$ in \mathcal{C} the following diagram commutes:

$$\begin{array}{ccccc} A & & F(A) & \xrightarrow{\eta_A} & G(A) \\ \downarrow f & & \downarrow F(f) & & \downarrow G(f) \\ B & & F(B) & \xrightarrow{\eta_B} & G(B) \end{array}$$

Natural transformation has, *nomen omen*, natural properties. One particularly interesting to us is the following fact.

Fact 2.20. Let η be a natural transformation of functors F, G from category \mathcal{C} to \mathcal{D} . Then η is an isomorphism if and only if all of the component morphisms are isomorphisms.

Proof. Suppose that η_A is an isomorphism for every $A \in \mathcal{C}$, where $\eta_A : F(A) \rightarrow G(A)$ is the morphism of the natural transformation corresponding to A . Then η^{-1} is simply given by the morphisms η_A^{-1} .

Now assume that η is an isomorphism, i.e. $\eta^{-1} \circ \eta = \text{id}_F$. *Ad contrario* assume that there is $A \in \mathcal{C}$ such that the component morphism $\eta_A : F(A) \rightarrow G(A)$ is not an isomorphism. It means that $\eta_A^{-1} \circ \eta_A \neq \text{id}_A$, hence $F(A) = \text{Dom}(\eta^{-1} \circ \eta)(A) \neq \text{Rng}(\eta^{-1} \circ \eta)(A) = F(A)$, which is obviously a contradiction. \square

Definition 2.21. In category theory, a *diagram* of type \mathcal{J} in category \mathcal{C} is a functor $D : \mathcal{J} \rightarrow \mathcal{C}$. \mathcal{J} is called the *index category* of D . In other words, D is of *shape* \mathcal{J} .

For example, $\mathcal{J} = \{-1 \leftarrow 0 \rightarrow 1\}$, then a diagram $D : \mathcal{J} \rightarrow \mathcal{C}$ is called a *cospan*. For example, if A, B, C are objects of \mathcal{C} and $f \in \text{Mor}(C, A), g \in \text{Mor}(C, B)$, then the following diagram is a cospan:

$$\begin{array}{ccc} A & & B \\ & \swarrow f & \nearrow g \\ & C & \end{array}$$

From now we omit explicit definition of the index category, as it is easily referable from a picture.

Definition 2.22. Let A, B, C, D be objects in the category \mathcal{C} with morphisms $e : C \rightarrow A, f : C \rightarrow B, g : A \rightarrow D, h : B \rightarrow D$ such that $g \circ e = h \circ f$. Then the following diagram:

$$\begin{array}{ccccc} & & D & & \\ & g \nearrow & & \nwarrow h & \\ A & & & & B \\ & e \nwarrow & & \nearrow f & \\ & & C & & \end{array}$$

is called a *pushout diagram*.

In both definitions of cospan and pushout diagrams we say that the object C is the *base* of the diagram.

Definition 2.23. The *cospan category* of category \mathcal{C} , referred to as $\text{Cospan}(\mathcal{C})$, is the category of cospan diagrams of \mathcal{C} , where morphisms between two cospans are natural transformations of the underlying functors.

We define *pushout category* analogously and call it $\text{Pushout}(\mathcal{C})$.

From now on we work in subcategories of cospan diagrams and pushout diagrams where we fix the base structure. Formally, for a fixed $C \in \mathcal{C}$, category $\text{Cospan}_C(\mathcal{C})$ is the category of all cospans in $\text{Cospan}(\mathcal{C})$ such that the base of the diagram is C . Natural transformation η of two diagrams in $\text{Cospan}_C(\mathcal{C})$ are such that the morphism $\eta_C : C \rightarrow C$ is an automorphism of C . $\text{Pushout}_C(\mathcal{C})$ is defined analogously. In most contexts we consider only one base structure, hence we will often write $\text{Pushout}(\mathcal{C})$ instead of $\text{Pushout}_C(\mathcal{C})$.

3. FRAÏSSÉ CLASSES

In this section we will take a closer look at classes of finitely generated structures with some characteristic properties. More specifically, we will describe a concept developed by a French mathematician Roland Fraïssé called Fraïssé limit.


3.1. Definitions.

Definition 3.1. Let L be a signature and M be an L -structure. The *age* of M is the class \mathcal{C} of all finitely generated structures that embed into M . The age of M is also associated with class of all finitely generated structures embeddable in M up to isomorphism.

Definition 3.2. We say that a class \mathcal{C} of finitely generated structures is *essentially countable* if it has countably many isomorphism types of finitely generated structures.

Definition 3.3. Let \mathcal{C} be a class of finitely generated structures. \mathcal{C} has the *hereditary property (HP)* if for any $A \in \mathcal{C}$ and any finitely generated substructure B of A it holds that $B \in \mathcal{C}$.

Definition 3.4. Let \mathcal{C} be a class of finitely generated structures. We say that \mathcal{C} has the *joint embedding property (JEP)* if for any $A, B \in \mathcal{C}$ there is a structure $C \in \mathcal{C}$ such that both A and B embed in C .



In terms of category theory we may say that \mathcal{C} is a category of finitely generated structures where morphisms are embeddings of those structures. Then the above diagram is a *span* diagram in category \mathcal{C} .


Fraïssé has shown fundamental theorems regarding age of a structure, one of them being the following one:

Fact 3.5. Suppose L is a signature and \mathcal{C} is a nonempty essentially countable set of finitely generated L -structures. Then \mathcal{C} has the HP and JEP if and only if \mathcal{C} is the age of some finite or countable structure.

Proof. One can read a proof of this fact in Wilfrid Hodges' classical book *Model Theory* [1, Theorem 7.1.1]. \square

Beside the HP and JEP Fraïssé has distinguished one more property of the class \mathcal{C} , namely the amalgamation property.

Definition 3.6. Let \mathcal{C} be a class of finitely generated L -structures. We say that \mathcal{C} has the *amalgamation property (AP)* if for any $A, B, C \in \mathcal{C}$ and embeddings $e: C \rightarrow A, f: C \rightarrow B$ there exists $D \in \mathcal{C}$ together with embeddings $g: A \rightarrow D$ and $h: B \rightarrow D$ such that $g \circ e = h \circ f$.



In terms of category theory, \mathcal{C} has the amalgamation property if every cospan diagram can be extended to a pushout diagram in category \mathcal{C} . We will get into more details later, in the definition of canonical amalgamation 3.19.

Definition 3.7. Class \mathcal{C} of finitely generated structures is a *Fraïssé class* if it is essentially countable, has HP, JEP and AP.


Definition 3.8. Let M be an L -structure. M is *ultrahomogeneous* if every isomorphism between finitely generated substructures of M extends to an automorphism of M .

Having those definitions we can provide the main Fraïssé theorem.

Theorem 3.9 (Fraïssé theorem). Let L be a countable language and let \mathcal{C} be a nonempty countable set of finitely generated L -structures which has HP, JEP and AP. Then \mathcal{C} is the age of a countable, ultrahomogeneous L -structure M . Moreover, M is unique up to isomorphism. We say that M is a Fraïssé limit of \mathcal{C} and denote this by $M = \text{Flim}(\mathcal{C})$.

Proof. Check the proof in [1, theorem 7.1.2]. \square

Definition 3.10. We say that an L -structure M is *weakly ultrahomogeneous* if for any A, B , finitely generated substructures of M , such that $A \subseteq B$ and an embedding $f: A \rightarrow M$ there is an embedding $g: B \rightarrow M$ which extends f .



Lemma 3.11. A countable structure is ultrahomogeneous if and only if it is weakly ultrahomogeneous.

Proof. Proof can be again found in [1, lemma 7.1.4(b)]. \square

This lemma will play a major role in the later parts of the paper. Weak ultrahomogeneity is an easier and more intuitive property and it will prove useful when recursively constructing the generic automorphism of a Fraïssé limit.

3.2. Random graph. In this section we'll take a closer look on a class of finite undirected graphs, which is a Fraïssé class.

The language of undirected graphs L consists of a single binary relational symbol E . If G is an L -structure then we call it a *graph*, and its elements *vertices*. If for some vertices $u, v \in G$ we have $G \models uEv$ then we say that there is an *edge* connecting u and v . If $G \models \forall x \forall y (xEy \leftrightarrow yEx)$ then we say that G is an *undirected graph*. From now on we consider only undirected graphs and omit the word *undirected*.

Proposition 3.12. *Let \mathcal{G} be the class of all finite graphs. \mathcal{G} is a Fraïssé class.*

Proof. \mathcal{G} is of course countable (up to isomorphism) and has the HP (substructure of a graph is also a graph). It has JEP: having two finite graphs G_1, G_2 take their disjoint union $G_1 \sqcup G_2$ as the extension of them both. \mathcal{G} has the AP. Having graphs A, B, C , where B and C are supergraphs of A , we can assume without loss of generality that $B \cap C = A$. Then $A \sqcup (B \setminus A) \sqcup (C \setminus A)$ is the graph we are looking for (with edges as in B and C and without any edges between $B \setminus A$ and $C \setminus A$). \square

Definition 3.13. The *random graph* is the Fraïssé limit of the class of finite graphs \mathcal{G} denoted by $\Gamma = \text{Flim}(\mathcal{G})$.

The concept of the random graph emerges independently in many fields of mathematics. For example, one can construct the graph by choosing at random for each pair of vertices if they should be connected or not. It turns out that the graph constructed this way is isomorphic to the random graph with probability 1.

The random graph Γ has one particular property that is unique to the random graph.

Fact 3.14 (Random graph property). *For each finite disjoint $X, Y \subseteq \Gamma$ there exists $v \in \Gamma \setminus (X \cup Y)$ such that $\forall u \in X$ we have that $\Gamma \models vEu$ and $\forall u \in Y$ we have that $\Gamma \models \neg vEu$.*

Proof. Take any finite disjoint $X, Y \subseteq \Gamma$. Let G_{XY} be the subgraph of Γ induced by the $X \cup Y$. Let $H = G_{XY} \cup \{w\}$, where w is a new vertex that does not appear in G_{XY} . Also, w is connected to all vertices of G_{XY} that come from X and to none of those that come from Y . This graph is of course finite, so it is embeddable in Γ by some $h: H \rightarrow \Gamma$. Let f be the partial isomorphism from $X \sqcup Y$ to $h[H] \subseteq \Gamma$, with X and Y projected to the part of $h[H]$ that come from X and Y respectively. By the ultrahomogeneity of Γ this isomorphism extends to an automorphism $\sigma \in \text{Aut}(\Gamma)$. Then $v = \sigma^{-1}(w)$ is the vertex we sought. \square

Fact 3.15. *If a countable graph G has the random graph property, then it is isomorphic to the random graph Γ .*

Proof. Enumerate vertices of both graphs: $\Gamma = \{a_1, a_2, \dots\}$ and $G = \{b_1, b_2, \dots\}$. We will construct a chain of partial isomorphisms $f_n: \Gamma \rightarrow G$ such that $\emptyset = f_0 \subseteq f_1 \subseteq f_2 \subseteq \dots$ and $a_n \in \text{Dom}(f_n)$ and $b_n \in \text{Rng}(f_n)$ for each $n \in \mathbb{N}$.

Suppose we have f_n . We seek $b \in G$ such that $f_n \cup \{(a_{n+1}, b)\}$ is a partial isomorphism. If $a_{n+1} \in \text{Dom } f_n$, then simply $b = f_n(a_{n+1})$. Otherwise, let $X = \{a \in \text{Dom } f_n \mid aE_\Gamma a_{n+1}\}$, $Y = \text{Dom } f_n \setminus X$, i.e. X are vertices of $\text{Dom } f_n$ that are connected with a_{n+1} in Γ and Y are those vertices that are not connected with a_{n+1} . Let b be a vertex of G that is connected to all vertices of $f_n[X]$ and to none $f_n[Y]$ (it exists by the random graph property). Then $f_n \cup \{(a_{n+1}, b)\}$ is a partial isomorphism. We find a for the b_{n+1} in the similar manner, so that $f_{n+1} = f_n \cup \{(a_{n+1}, b), (a, b_{n+1})\}$ is a partial isomorphism.

Finally, $f = \bigcup_{n=0}^{\infty} f_n$ is an isomorphism between Γ and G . Take any $a, b \in \Gamma$. Then for some big enough n we have that $aE_\Gamma b \Leftrightarrow f_n(a)E_G f_n(b) \Leftrightarrow f(a)E_G f(b)$. \square

Using this fact one can show that the graph constructed in the probabilistic manner is in fact isomorphic to the random graph Γ .

Definition 3.16. We say that a Fraïssé class \mathcal{C} has the *weak Hrushovski property (WHP)* if for every $A \in \mathcal{C}$ and an isomorphism of its finitely generated substructures $p: A \rightarrow A$ (also called a partial automorphism of A), there is some $B \in \mathcal{C}$ such that p can be extended to an automorphism of B , i.e. there is an embedding $i: A \rightarrow B$ and a $\bar{p} \in \text{Aut}(B)$ such that the following diagram commutes:


$$\begin{array}{ccc} B & \xrightarrow{\bar{p}} & B \\ \uparrow i & & \uparrow i \\ A & \xrightarrow{p} & A \end{array}$$

Proposition 3.17. *The class of finite graphs \mathcal{G} has the weak Hrushovski property.*

The proof of this proposition can be done directly, in a combinatorial manner, as shown in [8]. Hrushovski has also shown in [2] that finite graphs have stronger property, where each graph can be extended to a supergraph so that every partial automorphism of the graph extend to an automorphism of the supergraph.

Moreover, there is a theorem saying that every Fraïssé class \mathcal{C} , in a relational language L , with *free amalgamation* (see the definition 3.18 below) has WHP. The statement and proof of this theorem can be found in [9, theorem 3.2.8]. We provide the definition of free amalgamation that is coherent with the notions established in our paper.

Definition 3.18. Let L be a relational language and \mathcal{C} a class of L -structures. \mathcal{C} has *free amalgamation* if for every $A, B, C \in \mathcal{C}$ such that $C = A \cap B$ the following diagram commutes:




and $A \sqcup_C B \in \mathcal{C}$. $A \sqcup_C B$ here is an L -structure with domain $A \cup B$ such that for every n -ary symbol R from L , n -tuple $\bar{a} \subseteq A \cup B$, we have that $A \sqcup_C B \models R(\bar{a})$ if and only if $[\bar{a} \subseteq A$ and $A \models R(\bar{a})]$ or $[\bar{a} \subseteq B$ and $B \models R(\bar{a})]$.

Actually we did already implicitly work with free amalgamation in the Proposition 3.12, showing that the class of finite graphs is indeed a Fraïssé class.

3.3. Canonical amalgamation. Recall, $\text{Cospan}(\mathcal{C})$, $\text{Pushout}(\mathcal{C})$ are the categories of cospan and pushout diagrams of the category \mathcal{C} . We have also denoted the notion of cospans and pushouts with a fixed base structure C denoted as $\text{Cospan}_C(\mathcal{C})$ and $\text{Pushout}_C(\mathcal{C})$.


Definition 3.19. Let \mathcal{C} be a class of finitely generated L -structures. We say that \mathcal{C} has *canonical amalgamation* if for every $C \in \mathcal{C}$ there is a functor $\otimes_C: \text{Cospan}_C(\mathcal{C}) \rightarrow \text{Pushout}_C(\mathcal{C})$ with following properties:

- Let $A \leftarrow C \rightarrow B$ be a cospan. Then \otimes_C sends it to a pushout that preserves “the bottom” structures and embeddings, i.e.:



We have deliberately omitted names for embeddings of C . Of course, the functor has to take them into account, but this abuse of notation is convenient and should not lead into confusion.

- Let $A \leftarrow C \rightarrow B, A' \leftarrow C \rightarrow B'$ be cospans with a natural transformation η given by $\alpha: A \rightarrow A', \beta: B \rightarrow B', \gamma: C \rightarrow C$. Then \otimes_C preserves the morphisms of η when sending it to the natural transformation of pushouts by adding the $\delta: A \otimes_C B \rightarrow A' \otimes_C B'$ morphism:




Remark 3.20. *Free amalgamation is canonical.*

From now on in the paper, when A is an L -structure and α is an automorphism of A , then by (A, α) we mean the structure A expanded by the unary function corresponding to α , and A constantly denotes the L -structure.

Theorem 3.21. *Let \mathcal{C} be a Fraïssé class of L -structures with canonical amalgamation. Then the class \mathcal{D} of L -structures with automorphism is a Fraïssé class.*

Proof. \mathcal{D} is obviously countable and has HP. It suffices to show that it has AP (JEP follows by taking C to be the empty structure). Take any $(A, \alpha), (B, \beta), (C, \gamma) \in \mathcal{D}$ such that (C, γ) embeds into (A, α) and (B, β) . Then α, β, γ yield an automorphism η (as a natural transformation, see 2.20) of a cospan:



Then, by the Fact 2.18, $\otimes_C(\eta)$ is an automorphism of the pushout diagram that looks exactly like the diagram in the second point of the Definition 3.19. This means that the morphism $\delta: A \otimes_C B \rightarrow A' \otimes_C B'$ has to be automorphism. Thus, by the fact that the diagram commutes, we have the amalgamation of (A, α) and (B, β) over (C, γ) in \mathcal{D} . \square

The following theorem is one of the most important in construction of the generic automorphism given in the next section. Together with canonical amalgamation it gives a general fact about Fraïssé classes, namely it says that expanding a Fraïssé class with an automorphism of the structures does not change the limit.

Theorem 3.22. *Let \mathcal{C} be a Fraïssé class of finitely generated L -structures. Let \mathcal{D} be the class of structures from \mathcal{C} with additional unary function symbol interpreted as an automorphism of the structure. If \mathcal{C} has the weak Hrushovski property and \mathcal{D} is a Fraïssé class then the Fraïssé limit of \mathcal{C} is isomorphic to the Fraïssé limit of \mathcal{D} reduced to the language L .*

Proof. Let $\Gamma = \text{Flim}(\mathcal{C})$ and $(\Pi, \sigma) = \text{Flim}(\mathcal{D})$. By the Fraïssé Theorem 3.9 it suffices to show that the age of Π is \mathcal{C} and that it is weakly ultrahomogeneous. The former comes easily, as for every structure $A \in \mathcal{C}$ we have the structure $(A, \text{id}_A) \in \mathcal{D}$, which means that the structure A embeds into Π . On the other hand, if a structure $(B, \beta) \in \mathcal{D}$ embeds into (Π, σ) , then obviously $B \in \mathcal{C}$ by the definition of \mathcal{D} . Hence, \mathcal{C} is indeed the age of Π .

Now, to show that Π is weakly ultrahomogeneous, take any structures $A, B \in \mathcal{C}$ such that $A \subseteq B$ with a fixed embedding of A into Π . Without the loss of generality assume that $A = B \cap \Pi$ (i.e. A embeds into Π by inclusion). Let $\bar{A} \subseteq \Pi$ be the smallest substructure closed under the automorphism σ and containing A . It is finitely generated as an L -structure, as \mathcal{C} is the age of Π . Let C be a finitely generated structure such that $\bar{A} \rightarrow C \leftarrow B$. Such structure exists by the JEP of \mathcal{C} . Again, we may assume without the loss of generality that $\bar{A} \subseteq C$. Then $\sigma \upharpoonright_{\bar{A}}$ is a partial automorphism of C , hence by the WHP it can be extended to a structure $(\bar{C}, \gamma) \in \mathcal{D}$ such that $\gamma \upharpoonright_{\bar{A}} = \sigma \upharpoonright_{\bar{A}}$.

Then, by the weak ultrahomogeneity of (Π, σ) we can find an embedding g of (\bar{C}, γ) such that the following diagram commutes:

$$\begin{array}{ccc} (\bar{A}, \sigma \upharpoonright_{\bar{A}}) & \xrightarrow{\subseteq} & (\Pi, \sigma) \\ \downarrow \subseteq & \nearrow g & \\ (\bar{C}, \gamma) & & \end{array}$$

Thus, we have that the following diagram commutes:

$$\begin{array}{ccccc} A & \xrightarrow{\subseteq} & \bar{A} & \xrightarrow{\subseteq} & \Pi \\ \downarrow \subseteq & & \downarrow \subseteq & & \uparrow g \\ B & \xrightarrow{f} & C & \xrightarrow{\subseteq} & \bar{C} \end{array}$$

which proves that Π is indeed a weakly ultrahomogeneous structure. Hence, it is isomorphic to Γ . \square

Corollary 3.23. *Let \mathcal{C} be a Fraïssé class of finitely generated L -structures with WHP and canonical amalgamation. Let \mathcal{D} be the class consisting of structures from \mathcal{C} with an additional automorphism. Let $\Gamma = \text{Flim}(\mathcal{C})$ and $(\Pi, \sigma) = \text{Flim}(\mathcal{D})$. Then $\Gamma \cong \Pi$.*

Proof. It follows from Theorems 3.21 and 3.22. \square

4. CONJUGACY CLASSES IN AUTOMORPHISM GROUPS

Let M be a countable L -structure. Recall, we define a topology on the $G = \text{Aut}(M)$: for any finite function $f : M \rightarrow M$ we have a basic open set $[f]_G = \{g \in G \mid f \subseteq g\}$.

4.1. Prototype: pure set. In this section, $M = (M, =)$ is an infinite countable set (with no structure beyond equality).

Remark 4.1. If $f_1, f_2 \in \text{Aut}(M)$, then f_1 and f_2 are conjugate if and only if for each $n \in \mathbb{N} \cup \{\aleph_0\}$, f_1 and f_2 have the same number of orbits of size n .

Proof. It is easy to see. \square

Theorem 4.2. Let $\sigma \in \text{Aut}(M)$ be an automorphism with no infinite orbit and with infinitely many orbits of size n for every $n > 0$. Then the conjugacy class of σ is comeagre in $\text{Aut}(M)$.

Proof. We will show that the conjugacy class of σ is an intersection of countably many comeagre sets.

Let $A_n = \{\alpha \in \text{Aut}(M) \mid \alpha \text{ has infinitely many orbits of size } n\}$. This set is comeagre for every $n > 0$. Indeed, we can represent this set as an intersection of countable family of open dense sets. Let $B_{n,k}$ be the set of all finite functions $\beta: M \rightarrow M$ that consist of exactly k distinct n -cycles. Then:

$$\begin{aligned} A_n &= \{\alpha \in \text{Aut}(M) \mid \alpha \text{ has infinitely many orbits of size } n\} \\ &= \bigcap_{k=1}^{\infty} \{\alpha \in \text{Aut}(M) \mid \alpha \text{ has at least } k \text{ orbits of size } n\} \\ &= \bigcap_{k=1}^{\infty} \bigcup_{\beta \in B_{n,k}} [\beta]_{\text{Aut}(M)}, \end{aligned}$$

where indeed, $\bigcup_{\beta \in B_{n,k}} [\beta]_{\text{Aut}(M)}$ is dense in $\text{Aut}(M)$: take any finite $\gamma: M \rightarrow M$ such that $[\gamma]_{\text{Aut}(M)}$ is nonempty. Then also $\bigcup_{\beta \in B_{n,k}} [\beta]_{\text{Aut}(M)} \cap [\gamma]_{\text{Aut}(M)} \neq \emptyset$, one can easily construct a permutation that extends γ and has at least k many n -cycles.

Now we see that $A = \bigcap_{n=1}^{\infty} A_n$ is a comeagre set consisting of all functions that have infinitely many n -cycles for each n . The only thing left to show is that the set of functions with no infinite cycle is also comeagre. Indeed, for $m \in M$ let $B_m = \{\alpha \in \text{Aut}(M) \mid m \text{ has finite orbit in } \alpha\}$. This is an open dense set. It is a union over basic open sets generated by finite permutations with m in their domain. Denseness is also easy to see.

Finally, by the Remark 4.1, we can say that

$$\sigma^{\text{Aut}(M)} = \bigcap_{n=1}^{\infty} A_n \cap \bigcap_{m \in M} B_m,$$

which concludes the proof. \square

4.2. More general structures.

Fact 4.3. Suppose M is an arbitrary structure and $f_1, f_2 \in \text{Aut}(M)$. Then f_1 and f_2 are conjugate if and only if $(M, f_1) \cong (M, f_2)$ as structures with one additional unary function that is an automorphism.

Proof. Suppose that $f_1 = g^{-1}f_2g$ for some $g \in \text{Aut}(M)$. Then g is the isomorphism between (M, f_1) and (M, f_2) . On the other hand if $g: (M, f_1) \rightarrow (M, f_2)$ is an isomorphism, then $g \circ f_1 = f_2 \circ g$ which exactly means that f_1, f_2 conjugate. \square

Theorem 4.4. *Let \mathcal{C} be a Fraïssé class of finitely generated L -structures. Let \mathcal{D} be the class of structures from \mathcal{C} with additional unary function symbol interpreted as an automorphism of the structure. If \mathcal{C} has the weak Hrushovski property and \mathcal{D} is a Fraïssé class, then there is a comeagre conjugacy class in the automorphism group of the $\text{Flim}(\mathcal{C})$.*

Before we get to the proof, it is important to mention that an isomorphism between two finitely generated structures is uniquely given by a map from generators of one structure to the other. This allow us to treat a finite function as an isomorphism of finitely generated structures (if it yields one) and *vice versa*.

Proof. Let $\Gamma = \text{Flim}(\mathcal{C})$ and $(\Pi, \sigma) = \text{Flim}(\mathcal{D})$. First, by the Theorem 3.22, we may assume without the loss of generality that $\Pi = \Gamma$. Let $G = \text{Aut}(\Gamma)$, i.e. G is the automorphism group of Γ . We will construct a winning strategy for the second player in the Banach-Mazur game (see 2.7) on the topological space G with A being σ 's conjugacy class. By the Banach-Mazur theorem (see 2.13) this will prove that this class is comeagre.

Recall, G has a basis consisting of sets $\{g \in G \mid g \upharpoonright_A = g_0 \upharpoonright_A\}$ for some finite set $A \subseteq \Gamma$ and some automorphism $g_0 \in G$. In other words, a basic open set is a set of all extensions of some partial automorphism g_0 of finitely generated substructures of Γ . By $B_g \subseteq G$ we denote a basic open subset given by a partial isomorphism g . Again, Note that B_g is nonempty because of ultrahomogeneity of Γ .

With the use of Corollary 2.17 we can consider only games where both players choose partial isomorphisms. Namely, player I picks functions f_0, f_1, \dots and player II chooses g_0, g_1, \dots such that $f_0 \subseteq g_0 \subseteq f_1 \subseteq g_1 \subseteq \dots$, which identify the corresponding basic open subsets $B_{f_0} \supseteq B_{g_0} \supseteq \dots$

Our goal is to choose g_i in such a manner that $\bigcap_{i=0}^{\infty} B_{g_i} = \{g\}$ and (Γ, g) is ultrahomogeneous with age \mathcal{D} . By the Fraïssé theorem (see 3.9) it will follow that $(\Gamma, \sigma) \cong (\Gamma, g)$, thus by the Fact 4.3 we have that σ and g conjugate.

Fix a bijection $\gamma : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ such that for any $n, m \in \mathbb{N}$ we have $\gamma(n, m) \geq n$. This bijection naturally induces a well ordering on $\mathbb{N} \times \mathbb{N}$. This will prove useful later, as the main ingredient of the proof will be a bookkeeping argument.

For technical reasons, let $g_{-1} = \emptyset$ and $X_{-1} = \emptyset$. Enumerate the elements of the Fraïssé limit $\Gamma = \{v_0, v_1, \dots\}$. Suppose that player I in the n -th move chooses a partial automorphism f_n . We will construct a partial automorphism $g_n \supseteq f_n$ together with a finitely generated substructure $\Gamma_n \subseteq \Gamma$ and a set $X_n \subseteq \mathbb{N}^2$ such that the following properties hold:

- (i) g_n is a partial automorphism of Γ and an automorphism of finitely generated substructure Γ_n ,
- (ii) $g_n(v_n)$ and $g_n^{-1}(v_n)$ are defined.

Before we give the third point, suppose recursively that g_{n-1} already satisfy all those properties. Let us enumerate $\{(A_{n,k}, \alpha_{n,k}), (B_{n,k}, \beta_{n,k}), f_{n,k}\}_{k \in \mathbb{N}}$ all pairs of finitely generated structures with automorphisms such that the first substructure embed into the second by inclusion, i.e. $(A_{n,k}, \alpha_{n,k}) \subseteq (B_{n,k}, \beta_{n,k})$, and $f_{n,k}$ is an embedding of $(A_{n,k}, \alpha_{n,k})$ in the (Γ_{n-1}, g_{n-1}) . We allow $A_{n,k}$ to be empty. Although g_{n-1} is a finite function, we may treat it as a partial automorphism as we have said before.

- (iii) Let $(i, j) = \min\{(\{0, 1, \dots, n\} \times \mathbb{N}) \setminus X_{n-1}\}$ (with the order induced by γ). Then $X_n = X_{n-1} \cup \{(i, j)\}$ and $(B_{i,j}, \beta_{i,j})$ embeds in (Γ_n, g_n) so that this diagram commutes:

$$\begin{array}{ccc}
 & (\Gamma_n, g_n) & \\
 \hat{f}_{i,j} \nearrow & & \searrow \subseteq \\
 (B_{i,j}, \beta_{i,j}) & & (\Gamma_{n-1}, g_{n-1}) \\
 \subseteq \nwarrow & & \nearrow f_{i,j} \\
 & (A_{i,j}, \alpha_{i,j}) &
 \end{array}$$

First, we will satisfy the item (iii). Namely, we will construct Γ'_n, g'_n such that $g_{n-1} \subseteq g'_n$, $\Gamma_{n-1} \subseteq \Gamma'_n$, g'_n gives an automorphism of Γ'_n and $f_{i,j}$ extends to an embedding of $(B_{i,j}, \beta_{i,j})$ to (Γ'_n, g'_n) . But this can be easily done by the fact, that \mathcal{D} has the amalgamation property.

It is important to note that g'_n should be a finite function and once again, as it is an automorphism of a finitely generated structure, we may think it is simply a map from one generators of Γ'_n to the others. By the weak ultrahomogeneity of Γ , we may assume that $\Gamma'_n \subseteq \Gamma$.

Now, by the WHP of \mathcal{C} we can extend $\langle \Gamma'_n \cup \{v_n\} \rangle$ together with its partial isomorphism g'_n to a finitely generated structure Γ_n together with its automorphism $g_n \supseteq g'_n$ and (again by weak ultrahomogeneity) without the loss of generality we may assume that $\Gamma_n \subseteq \Gamma$. This way we've constructed g_n that has all desired properties.

Now we need to see that $g = \bigcap_{n=0}^{\infty} g_n$ is indeed an automorphism of Γ such that (Γ, g) has the age \mathcal{D} and is weakly ultrahomogeneous. It is of course an automorphism of Γ as it is defined for every $v \in \Gamma$ and is an union of an increasing chain of automorphisms of finitely generated substructures.

Take any $(B, \beta) \in \mathcal{D}$. Then, there are i, j such that $(B, \beta) = (B_{i,j}, \beta_{i,j})$ and $A_{i,j} = \emptyset$. By the bookkeeping there was n such that $(i, j) = \min\{\{0, 1, \dots, n\} \times \mathbb{N} \setminus X_n\}$. This means that (B, β) embeds into (Γ_n, g_n) , hence it embeds into (Γ, g) . Thus, \mathcal{D} is a subclass of the age of (Γ, g) . The other inclusion is obvious. Hence, the age of (Γ, g) is \mathcal{H} .

It is also weakly ultrahomogeneous. Having $(A, \alpha) \subseteq (B, \beta)$, and an embedding $f : (A, \alpha) \rightarrow (\Gamma, g)$, we may find $n \in \mathbb{N}$ such that $(i, j) = \min\{\{0, 1, \dots, n-1\} \times X_{n-1}\}$ and $(A, \alpha) = (A_{i,j}, \alpha_{i,j})$, $(B, \beta) = (B_{i,j}, \beta_{i,j})$ and $f = f_{i,j}$. This means that there is a compatible embedding of (B, β) into (Γ_n, g_n) , which means we can also embed it into (Γ, g) .

Hence, $(\Gamma, g) \cong (\Gamma, \sigma)$. By this we know that g and σ are conjugate in G , thus player II have a winning strategy in the Banach-Mazur game with $A = \sigma^G$, thus σ^G is comeagre in G and σ is a generic automorphism. \square

Theorem 4.5. *Let \mathcal{C} be a Fraïssé class of finitely generated L -structures with WHP and canonical amalgamation. Then $\text{Flim}(\mathcal{C})$ has a generic automorphism.*

Proof. It follows trivially from Corollary 3.23 and the above Theorem 4.4. \square

4.3. Properties of the generic automorphism. Let \mathcal{C} be a Fraïssé class of finitely generated L -structures with weak Hrushovski property and canonical

amalgamation. Let \mathcal{D} be the Fraïssé class (by the Theorem 4.5 of the structures of \mathcal{C} with additional automorphism of the structure. Let $\Gamma = \text{Flim}(\mathcal{C})$.

Proposition 4.6. *Let σ be the generic automorphism of Γ . Then the set of fixed points of σ is isomorphic to Γ .*

Proof. Let $S = \{x \in \Gamma \mid \sigma(x) = x\}$. First we need to show that it is an infinite. By the theorem 4.4 we know that (Γ, σ) is the Fraïssé limit of \mathcal{D} , thus we can embed finite L -structures of any size with identity as an automorphism of the structure into (Γ, σ) . Thus S has to be infinite. Also, the same argument shows that the age of the structure is exactly \mathcal{C} . It is weakly ultrahomogeneous, also by the fact that (Γ, σ) is in \mathcal{D} . \square

5. EXAMPLES

Example 5.1. Let \mathcal{L} be the class of all finite linear orderings. Then:

- (1) \mathcal{L} is a Fraïssé class.
- (2) \mathcal{L} has canonical amalgamation.
- (3) \mathcal{L} does not have WHP.

\mathcal{L} of course has HP and is essentially countable. JEP is also easy, as having two finite linear orderings we can just embed the one with fewer elements into the bigger one.

We will show that \mathcal{L} has canonical amalgamation (CAP). Let C be a finite linear ordered set. We will define \otimes_C . Let A, B be finite linear orderings that C embeds into. We may suppose that $C = A \cap B$. Then we define an ordering on $D = A \cup B$. For $d, e \in D$, let $d \leq_D e$ if one of the following hold:

- $d, e \in A$ and $d \leq_A e$,
- $d, e \in B$ and $d \leq_B e$,
- $d \in A, e \in B$ and there is $c \in C$ such that $d \leq_A c$ and $c \leq_B e$,
- $d \in B, e \in A$ and there is $c \in C$ such that $d \leq_B c$ and $c \leq_A e$,
- $d \in A, e \in B$ and for all $c \in C$ $d \leq_A c \Leftrightarrow e \leq_B c$

One can imagine that D can be constructed by laying elements of C in a row and putting elements of A and B appropriately between elements of C with all elements of A to the left and all elements of B to the right between two adjacent elements of C . This clearly is a canonical amalgamation.





FIGURE 1. Visual representation of the construction.

On the other hand \mathcal{L} cannot have *WHP*. This follows from the fact that the only automorphism of a finite linear ordering is identity, so we cannot extend a partial automorphism sending exactly one element to some distinct element. However, in this case, generic automorphism exists which was shown by Truss [10].

Definition 5.2. Let X be a set. A ternary relation $\leq^C \subseteq X^3$ is a *cyclic order*, where we denote $(a, b, c) \in \leq^C$ as $a \leq_b^C c$ (or simply $a \leq_b c$ when there's only one relation in the context), when it satisfies the following properties:

- If $a \leq_b c$, then $b \leq_c a$.
- If $a \leq_b c$, then not $c \leq_b a$.
- If $a \leq_b c$ and $a \leq_c d$, then $a \leq_b d$.
- If a, b, c are pairwise distinct, then either $a \leq_b c$ or $c \leq_b a$.

It is easy to visualize a cyclic ordering as a directed (*nomen omen*) cycle. For example, a 11-element cyclic order could be drawn like this:



For three elements a, b, c we say that $a \leq_b c$ if after "cutting" the cycle at b we have a path from a to c . In this particular example we can say that the green element is red-element-smaller than the blue one.

Example 5.3. The class \mathcal{C} of all finite cyclic orders is a Fraïssé class, but does not have *WHP* or *CAP*.

It is not hard to show that \mathcal{C} is indeed the Fraïssé class. As usual, the hardest part is showing *AP*, which in this case is done analogously to the linear orders. The Fraïssé limit of \mathcal{C} is a countable unit circle.

\mathcal{C} hasn't *WHP* by the similar argument to this for linear orderings. Imagine a cycling order of three elements and a partial automorphism with one fixed point and moving second element to the third. This cannot be extended to automorphism of any finite cyclic order.

Also, \mathcal{C} cannot have *CAP*. A reason to that is that it do not admit canonical amalgamation over the empty structure (see this by taking 1-element cyclic order and 3-element cyclic order with automorphism other than identity).

In contrast to linear orderings the Fraïssé limit $\Sigma = \text{Flim } \mathcal{C}$ has no generic automorphism. Consider the set A of automorphisms of Σ with at least one

finite orbit of size greater than 1. It is open, not dense and closed on conjugation. Openness follows from the fact that all finite orbits of a given automorphism have the same size. Thus A can be represented as a union of basic set generated by finite cycles of length greater than 1. It is not dense, as it has empty intersection with basic set generated by identity of a single element. It is also closed on taking conjugation, as the order of elements does not change when conjugating. Thus there cannot be a dense conjugacy class in $\text{Aut}(\Sigma)$ and so there's no generic automorphism.

Example 5.4. The class \mathcal{V} of all finitely generated vector spaces over a countable field is a Fraïssé class with WHP and CAP.

The prove of this is relatively easy, knowing that there is essentially one vector space of every finite dimension and that every linear independent subset of a vector space can be extended to a basis of this space. The Fraïssé limit of \mathcal{V} is the ω -dimensional vector space. Thus, by our key Theorem 4.5 we know that it has a generic automorphism.

Example 5.5. The class of all finite graphs \mathcal{G} is a Fraïssé class with WHP and free amalgamation.

We have already shown this fact. Thus get that the random graph has a generic automorphism.

Example 5.6. A K_n -free graph is a graph with no n -clique as its subgraph. Let \mathcal{G}_n be the class of finite K_n -free graphs. \mathcal{G}_n is a Fraïssé class with WHP and free amalgamation.

Showing that \mathcal{G}_n is indeed a Fraïssé class is almost the same as in normal graphs, together with free amalgamation. WHP is trickier and the proof can be seen in [7] Theorem 3.6. Hence, $\text{Flim}(\mathcal{G}_n)$ has a generic automorphism.

REFERENCES

- [1] Wilfrid Hodges. *Model Theory*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1993. DOI: [10.1017/CB09780511551574](https://doi.org/10.1017/CB09780511551574).
- [2] Ehud Hrushovski. “Extending partial isomorphisms of graphs”. In: *Combinatorica* 12 (1992), pp. 411–416. DOI: <https://doi.org/10.1007/BF01305233>.
- [3] A. A. Ivanov. “Generic expansions of ω -categorical structures and semantics of generalized quantifiers”. In: *Journal of Symbolic Logic* 64.2 (1999). DOI: [10.2307/2586500](https://doi.org/10.2307/2586500).
- [4] Alexander S. Kechris. *Classical Descriptive Set Theory*. Springer New York, NY, 1995. DOI: [10.1007/978-1-4612-4190-4](https://doi.org/10.1007/978-1-4612-4190-4).
- [5] Alexander S. Kechris and Christian Rosendal. “Turbulence, amalgamation, and generic automorphisms of homogeneous structures”. In: *Proceedings of the London Mathematical Society* 94.2 (2007), pp. 302–350. DOI: <https://doi.org/10.1112/plms/pdl007>.
- [6] Saunders Mac Lane. *Categories for the Working Mathematician*. Springer New York, NY, 1978. DOI: [10.1007/978-1-4757-4721-8](https://doi.org/10.1007/978-1-4757-4721-8).

- [7] Shixiao Liu. “Fraïssé Limits, Hrushovski Property and Generic Automorphisms”. In: (2017). URL: https://logic.pku.edu.cn/ann_attachments/eppa-presentation.pdf.
- [8] Cédric Milliet. “Extending partial isomorphisms of finite graphs”. In: (2004). URL: <http://math.univ-lyon1.fr/~milliet/grapheanglais.pdf>.
- [9] Daoud Nasri Siniora. *Automorphism Groups of Homogeneous Structures*. The University of Leeds (Department of Pure Mathematics), 2017. URL: <https://core.ac.uk/download/pdf/83934818.pdf>.
- [10] J. K. Truss. “Generic Automorphisms of Homogeneous Structures”. In: *Proceedings of the London Mathematical Society* s3-65.1 (1992), pp. 121–141. DOI: <https://doi.org/10.1112/plms/s3-65.1.121>.